

# Model completeness of finitely ramified henselian valued field with various value groups

Anna De Mase

Università degli Studi della Campania "Luigi Vanvitelli"

7th December 2023

# Index

## 1 Valued fields

## 2 Transfer principles

- Model completeness for hens  $vf(0,p)$  valued in a  $Z$ -group
- Model completeness for hens  $vf(0,p)$  valued in an oag with finite spines
- Model completeness for hens  $vf(0,p)$  valued in an oag with spine of order type  $\omega^*$  and no colors



# Valued fields

Let  $K$  be a field and  $G$  a totally ordered abelian group. A *valuation* over  $K$  is a surjective map

$$v : K \longrightarrow G \cup \{\infty\}$$

satisfying the following conditions:

- $v(a) = \infty \iff a = 0$ ;
- $v(ab) = v(a) + v(b)$ ;
- $v(a + b) \geq \min\{v(a), v(b)\}$ .

$O = \{x \in K \mid v(x) \geq 0\}$  is the valuation ring. It is a local ring with unique maximal ideal  $M = \{x \in K \mid v(x) > 0\}$ . The quotient  $O/M = k$  is the *residue field* of the valuation  $v$ .

Cases:

- 1.  $\text{char } K = \text{char } k = 0$ ;
- 2.  $\text{char } K = 0, \text{char } k = p$ ;
- 3.  $\text{char } K = \text{char } k = p$ .



# Valued fields

Let  $K$  be a field and  $G$  a totally ordered abelian group. A *valuation* over  $K$  is a surjective map

$$v : K \longrightarrow G \cup \{\infty\}$$

satisfying the following conditions:

- $v(a) = \infty \iff a = 0$ ;
- $v(ab) = v(a) + v(b)$ ;
- $v(a + b) \geq \min\{v(a), v(b)\}$ .

$O = \{x \in K \mid v(x) \geq 0\}$  is the valuation ring. It is a local ring with unique maximal ideal  $M = \{x \in K \mid v(x) > 0\}$ . The quotient  $O/M = k$  is the *residue field* of the valuation  $v$ .

Cases:

- 1.  $\text{char } K = \text{char } k = 0$ ;
- 2.  $\text{char } K = 0, \text{char } k = p$ ;
- 3.  $\text{char } K = \text{char } k = p$ .



# Examples

1. Let  $p$  be a prime. The field of  $p$ -adic numbers

$$\mathbb{Q}_p = \left\{ a = \sum_{i \geq k} a_i p^i \mid k \in \mathbb{Z}, a_i \in \{0, \dots, p-1\} \right\}$$

with the valuation  $v_p(a) = \min\{i \mid a_i \neq 0\}$  (the  $p$ -adic valuation) with values in  $\mathbb{Z}$  and residue field  $\mathbb{F}_p$ .

2. Let  $k$  be a field and  $G$  an ordered abelian group. The valued field of *generalized power series* (or *Hahn field*)

$$k((G)) = \left\{ a = \sum_{g \in G} a_g t^g \mid a_g \in k, \text{ for all } g \in G \text{ and } \text{supp}(a) \text{ is well ordered} \right\}$$

with the valuation  $v_t(a) = \min\{g \mid a_g \neq 0\}$  (the  $t$ -adic valuation) with values in  $G$  and residue field  $k$ .



# Examples

1. Let  $p$  be a prime. The field of  $p$ -adic numbers

$$\mathbb{Q}_p = \left\{ a = \sum_{i \geq k} a_i p^i \mid k \in \mathbb{Z}, a_i \in \{0, \dots, p-1\} \right\}$$

with the valuation  $v_p(a) = \min\{i \mid a_i \neq 0\}$  (the  $p$ -adic valuation) with values in  $\mathbb{Z}$  and residue field  $\mathbb{F}_p$ .

2. Let  $k$  be a field and  $G$  an ordered abelian group. The valued field of *generalized power series* (or *Hahn field*)

$$k((G)) = \left\{ a = \sum_{g \in G} a_g t^g \mid a_g \in k, \text{ for all } g \in G \text{ and } \text{supp}(a) \text{ is well ordered} \right\}$$

with the valuation  $v_t(a) = \min\{g \mid a_g \neq 0\}$  (the  $t$ -adic valuation) with values in  $G$  and residue field  $k$ .



# Valued fields

## Definition

*A valued field is henselian if its valuation extends uniquely to every algebraic extension.*

**Remark.**  $(\mathbb{Q}_p, v_p)$  is henselian.

## Theorem (Hensel's Lemma)

*Suppose  $K$  is a pseudo-complete valued field, then it is henselian.*

## Definition

*Let  $K$  be a valued field. We say that  $K$  is finitely ramified if  $\text{char}(k) = p$  and  $|\{v(x) : 0 < v(x) \leq v(p)\}| = e < \infty$ . In particular, if  $e = 1$  the field is unramified. The element with minimal positive valuation is called uniformizer.*

## AKE - equicharacteristic zero

Consider  $\mathcal{L}_{rings} = \{+, \cdot, 0, 1\}$ ,  $\mathcal{L}_{oags} = \{+, 0, \leq\}$ , and  $\mathcal{L}_{vf} = (\mathcal{L}_{Rings}, \mathcal{L}_{oags}, \mathcal{L}_{rings}, v, res)$ .

## Theorem (Ax-Kochen-Ershov principle)

*Let  $(K_1, v_1), (K_2, v_2)$  be two henselian valued field whose residue fields  $k_1, k_2$  have characteristic 0 and let  $G_1, G_2$  be their value groups. Then*

$$K_1 \equiv_{vf} K_2 \text{ iff } k_1 \equiv_{rings} k_2 \text{ and } G_1 \equiv_{oag} G_2$$

Theorem (Ax-Kochen-Ershov principle ( $\preceq$ -version))

*Let  $(K_1, v_1), (K_2, v_2)$  be two henselian valued field whose residue fields  $k_1, k_2$  have characteristic 0 and let  $G_1, G_2$  be their value groups. Then*

$$K_1 \preceq_{vf} K_2 \text{ iff } k_1 \preceq_{rings} k_2 \text{ and } G_1 \preceq_{oag} G_2$$





# AKE - mixed characteristic

## Theorem (Bélair)

Let  $(K_1, v_1), (K_2, v_2)$  be two unramified henselian valued fields with perfect residue fields  $k_1, k_2$  and let  $G_1, G_2$  be their value groups. Then

$$K_1 \preceq_{vf} K_2 \text{ iff } k_1 \preceq_{rings} k_2 \text{ and } G_1 \preceq_{oag} G_2$$

## Theorem (Anscombe-Jahnke)

Let  $(K_1, v_1), (K_2, v_2)$  be two unramified henselian valued fields with arbitrary residue fields  $k_1, k_2$  and let  $G_1, G_2$  be their value groups. Then

$$K_1 \preceq_{vf} K_2 \text{ iff } k_1 \preceq_{rings} k_2 \text{ and } G_1 \preceq_{oag} G_2$$

**Question:** Does the transfer hold for valued fields with (finite) ramification? **Answer:** In general, no.



# Index

## 1 Valued fields

## 2 Transfer principles

- Model completeness for hens vf  $(0,p)$  valued in a  $Z$ -group
- Model completeness for hens vf  $(0,p)$  valued in an oag with finite spines
- Model completeness for hens vf  $(0,p)$  valued in an oag with spine of order type  $\omega^*$  and no colors



# $\mathbb{Z}$ -groups

## Definition

*An ordered abelian group is called a  $\mathbb{Z}$ -group if it is elementarily equivalent to  $\mathbb{Z}$  as an ordered abelian group.*

## Proposition

*The theory of  $\mathbb{Z}$  as an ordered abelian group has quantifier elimination in the Presburger language  $\mathcal{L}_{pres} = \{+, 0, 1, \leq, \equiv_m\}_{m \in \mathbb{N}}$ , where 1 is a constant for the minimal positive element and  $a \equiv_m b$  iff  $a - b \in m\mathbb{Z}$ .*



### Theorem (Derakhshan-Macintyre)

*Let  $K$  be a Henselian valued field of mixed characteristic  $(0, p)$  with finite ramification  $e \geq 1$ . Suppose the value group of  $K$  is a  $\mathbb{Z}$ -group. If the theory of the residue field  $k$  is model complete in the language of rings, then the theory of  $K$  is model complete in the language of rings.*



# Some observations

- (i) let  $k$  be a field. If  $Th(k)$  is model complete in  $\mathfrak{L}_{rings}$ , then  $k$  is perfect.
- (ii) let  $G$  be a  $\mathbb{Z}$ -group, then  $G/\mathbb{Z}$  is a divisible ordered abelian group.
- (iii) Let  $K$  be an henselian valued field of mixed characteristic  $(0, p)$  and ramification index  $e$ , and let  $n > e$  be an integer coprime with  $p$ . Then the valuation ring is existentially definable by the formula

$$\exists y(1 + px^n = y^n),$$

and the maximal ideal is existentially definable by the formula

$$\exists y(1 + \frac{1}{p}x^n = y^n).$$



# Some observations

- (i) let  $k$  be a field. If  $Th(k)$  is model complete in  $\mathcal{L}_{rings}$ , then  $k$  is perfect.
- (ii) let  $G$  be a  $\mathbb{Z}$ -group, then  $G/\mathbb{Z}$  is a divisible ordered abelian group.
- (iii) Let  $K$  be an henselian valued field of mixed characteristic  $(0, p)$  and ramification index  $e$ , and let  $n > e$  be an integer coprime with  $p$ . Then the valuation ring is existentially definable by the formula

$$\exists y(1 + px^n = y^n),$$

and the maximal ideal is existentially definable by the formula

$$\exists y(1 + \frac{1}{p}x^n = y^n).$$



# Some observations

- (i) let  $k$  be a field. If  $Th(k)$  is model complete in  $\mathcal{L}_{rings}$ , then  $k$  is perfect.
- (ii) let  $G$  be a  $\mathbb{Z}$ -group, then  $G/\mathbb{Z}$  is a divisible ordered abelian group.
- (iii) Let  $K$  be an henselian valued field of mixed characteristic  $(0, p)$  and ramification index  $e$ , and let  $n > e$  be an integer coprime with  $p$ . Then the valuation ring is existentially definable by the formula

$$\exists y(1 + px^n = y^n),$$

and the maximal ideal is existentially definable by the formula

$$\exists y(1 + \frac{1}{p}x^n = y^n).$$



# Sketch of the proof

Let  $(K_1, v_1), (K_2, v_2) \models Th(K, v)$  such that  $K_1 \subseteq K_2$ . Claim:  
 $(K_1, v_1) \preceq (K_2, v_2)$  in  $\mathcal{L}_{rings}$ .

Step 1. Assume  $K_1, K_2$   $\aleph_1$ -saturated;

Step 2. By (iii),  $(K_1, v_1) \subseteq (K_2, v_2)$ ;

Step 3. Coarsening and reduction to the equicharacteristic 0 case: by AKE and (ii) if  $\dot{K}_1 \preceq \dot{K}_2$ , then  $\dot{K}_1 \preceq \dot{K}_2$ ;

Step 4. By Bèlair's theorem, if  $k_1 \preceq k_2$  then  $W(k_1) \preceq W(k_2)$ , where  $W(k_i)$  is the Witt ring of  $k_i$ ,  $i = 1, 2$ .

Step 5. For  $i = 1, 2$  and an uniformizer  $\pi \in K_1 \subseteq K_2$ , interpret  $\dot{K}_i := W(k_i)(\pi)$  into  $W(k_i)$ . Thus  $K_1 \preceq K_2$  as fields. By (iii),  $K_1 \preceq K_2$  as valued fields in  $\mathcal{L}_{rings}$ .





# Sketch of the proof

Let  $(K_1, v_1), (K_2, v_2) \models Th(K, v)$  such that  $K_1 \subseteq K_2$ . Claim:  
 $(K_1, v_1) \preceq (K_2, v_2)$  in  $\mathcal{L}_{rings}$ .

Step 1. Assume  $K_1, K_2$   $\aleph_1$ -saturated;

Step 2. By (iii),  $(K_1, v_1) \subseteq (K_2, v_2)$ ;

Step 3. Coarsening and reduction to the equicharacteristic 0 case: by AKE and (ii) if  $\dot{K}_1 \preceq \dot{K}_2$ , then  $\dot{K}_1 \preceq \dot{K}_2$ ;

Step 4. By Bèlair's theorem, if  $k_1 \preceq k_2$  then  $W(k_1) \preceq W(k_2)$ , where  $W(k_i)$  is the Witt ring of  $k_i$ ,  $i = 1, 2$ .

Step 5. For  $i = 1, 2$  and an uniformizer  $\pi \in K_1 \subseteq K_2$ , interpret  $\dot{K}_i := W(k_i)(\pi)$  into  $W(k_i)$ . Thus  $K_1 \preceq K_2$  as fields. By (iii),  $K_1 \preceq K_2$  as valued fields in  $\mathcal{L}_{rings}$ .



# Sketch of the proof

Let  $(K_1, v_1), (K_2, v_2) \models Th(K, v)$  such that  $K_1 \subseteq K_2$ . Claim:  
 $(K_1, v_1) \preceq (K_2, v_2)$  in  $\mathcal{L}_{rings}$ .

Step 1. Assume  $K_1, K_2$   $\aleph_1$ -saturated;

Step 2. By (iii),  $(K_1, v_1) \subseteq (K_2, v_2)$ ;

Step 3. Coarsening and reduction to the equicharacteristic 0 case: by AKE and (ii) if  $\dot{K}_1 \preceq \dot{K}_2$ , then  $\dot{K}_1 \preceq \dot{K}_2$ ;

Step 4. By Bèlair's theorem, if  $k_1 \preceq k_2$  then  $W(k_1) \preceq W(k_2)$ , where  $W(k_i)$  is the Witt ring of  $k_i$ ,  $i = 1, 2$ .

Step 5. For  $i = 1, 2$  and an uniformizer  $\pi \in K_1 \subseteq K_2$ , interpret  $\dot{K}_i := W(k_i)(\pi)$  into  $W(k_i)$ . Thus  $K_1 \preceq K_2$  as fields. By (iii),  $K_1 \preceq K_2$  as valued fields in  $\mathcal{L}_{rings}$ .



# Sketch of the proof

Let  $(K_1, v_1), (K_2, v_2) \models Th(K, v)$  such that  $K_1 \subseteq K_2$ . Claim:  
 $(K_1, v_1) \preceq (K_2, v_2)$  in  $\mathcal{L}_{rings}$ .

Step 1. Assume  $K_1, K_2$   $\aleph_1$ -saturated;

Step 2. By (iii),  $(K_1, v_1) \subseteq (K_2, v_2)$ ;

Step 3. Coarsening and reduction to the equicharacteristic 0 case: by AKE and (ii) if  $\dot{K}_1 \preceq \dot{K}_2$ , then  $K_1 \preceq K_2$ ;

Step 4. By Bèlair's theorem, if  $k_1 \preceq k_2$  then  $W(k_1) \preceq W(k_2)$ , where  $W(k_i)$  is the Witt ring of  $k_i$ ,  $i = 1, 2$ .

Step 5. For  $i = 1, 2$  and an uniformizer  $\pi \in K_1 \subseteq K_2$ , interpret  $\dot{K}_i := W(k_i)(\pi)$  into  $W(k_i)$ . Thus  $K_1 \preceq K_2$  as fields. By (iii),  $K_1 \preceq K_2$  as valued fields in  $\mathcal{L}_{rings}$ .



# Sketch of the proof

Let  $(K_1, v_1), (K_2, v_2) \models Th(K, v)$  such that  $K_1 \subseteq K_2$ . Claim:  
 $(K_1, v_1) \preceq (K_2, v_2)$  in  $\mathcal{L}_{rings}$ .

Step 1. Assume  $K_1, K_2$   $\aleph_1$ -saturated;

Step 2. By (iii),  $(K_1, v_1) \subseteq (K_2, v_2)$ ;

Step 3. Coarsening and reduction to the equicharacteristic 0 case: by AKE and (ii) if  $\dot{K}_1 \preceq \dot{K}_2$ , then  $\dot{K}_1 \preceq \dot{K}_2$ ;

Step 4. By Bèlair's theorem, if  $k_1 \preceq k_2$  then  $W(k_1) \preceq W(k_2)$ , where  $W(k_i)$  is the Witt ring of  $k_i$ ,  $i = 1, 2$ .

Step 5. For  $i = 1, 2$  and an uniformizer  $\pi \in K_1 \subseteq K_2$ , interpret  $\dot{K}_i := W(k_i)(\pi)$  into  $W(k_i)$ . Thus  $K_1 \preceq K_2$  as fields. By (iii),  $K_1 \preceq K_2$  as valued fields in  $\mathcal{L}_{rings}$ .



# Sketch of the proof

Let  $(K_1, v_1), (K_2, v_2) \models Th(K, v)$  such that  $K_1 \subseteq K_2$ . Claim:  
 $(K_1, v_1) \preceq (K_2, v_2)$  in  $\mathcal{L}_{rings}$ .

Step 1. Assume  $K_1, K_2$   $\aleph_1$ -saturated;

Step 2. By (iii),  $(K_1, v_1) \subseteq (K_2, v_2)$ ;

Step 3. Coarsening and reduction to the equicharacteristic 0 case: by AKE and (ii) if  $\dot{K}_1 \preceq \dot{K}_2$ , then  $\dot{K}_1 \preceq \dot{K}_2$ ;

Step 4. By Bèlair's theorem, if  $k_1 \preceq k_2$  then  $W(k_1) \preceq W(k_2)$ , where  $W(k_i)$  is the Witt ring of  $k_i$ ,  $i = 1, 2$ .

Step 5. For  $i = 1, 2$  and an uniformizer  $\pi \in K_1 \subseteq K_2$ , interpret  $\dot{K}_i := W(k_i)(\pi)$  into  $W(k_i)$ . Thus  $K_1 \preceq K_2$  as fields. By (iii),  $K_1 \preceq K_2$  as valued fields in  $\mathcal{L}_{rings}$ .



# Index

## 1 Valued fields

## 2 Transfer principles

- Model completeness for hens vf  $(0,p)$  valued in a  $Z$ -group
- Model completeness for hens vf  $(0,p)$  valued in an oag with finite spines
- Model completeness for hens vf  $(0,p)$  valued in an oag with spine of order type  $\omega^*$  and no colors



# Oags with finite spines

Let  $G$  be an OAG. For each positive integer  $n$  we recall the definition of the spine  $S_n$ :

## Definition

For  $n \in \mathbb{N}$  and  $a \in G \setminus nG$ , let  $H_a$  be the largest convex subgroup of  $G$  such that  $a \notin H_a + nG$ ; set  $H_a = 0$  if  $a \in nG$ . Define  $S_n := G / \sim$ , with  $a \sim a'$  iff  $H_a = H_{a'}$ . and let  $s_n : G \rightarrow S_n$  be the canonical projection. For  $\alpha = s_n(a) \in S_n$ , define  $\overline{H_\alpha} := H_a$ . Since the system of convex subgroups of an ordered abelian group are linearly ordered,  $S_n$  is an interpretable set linearly ordered by  $\alpha \leq \alpha'$  if  $\overline{H_\alpha} \subseteq \overline{H_{\alpha'}}$ . The structure  $(S_n, <)$  is called the  $n$ -spine of  $G$ .



# Oags with finite spines

## Definition

*If  $G$  is an OAG such that for each  $n \in \mathbb{N}$ ,  $|S_n|$  is finite,  $G$  is said to have finite spines.*

Remark. All the  $\overline{H_\alpha}$  are definable in  $\mathcal{L}_{oag}$  and, moreover, if  $G$  is a group with finite spines, then  $\{\overline{H_\alpha} \mid \alpha \in S_n, n \in \mathbb{N}\}$  are all the definable convex subgroups of  $G$ .

$\implies G$  has only finitely or countably many definable convex subgroups that we denote with  $(H_i)_{i \in I}$ , where  $I$  is a finite or countable set of indexes.





## QE

## Proposition (Halevi-Hasson/Farré)

Let  $G$  be an ordered abelian group with finite spines and let  $\{H_i\}_{i \in I}$  be its definable convex subgroups for some  $I$  finite or countable. Then  $G$  has quantifier elimination in the language:

$$\mathcal{L} = \mathcal{L}_{\text{oag}} \cup \{(x =_{H_i} y + j_{G/H_i})_{i \in I, j \in \mathbb{N}}, (x \equiv_m^{H_i} y + j_{G/H_i})_{i \in I, j \in \mathbb{Z}, m \in \mathbb{N}}\}$$

where  $j_{G/H_i}$  is  $j$  times the minimal positive element of the quotient  $G/H_i$ , if it exists, 0 otherwise.



# Model completeness

## Proposition (D.M.)

*Let  $G$  be an ordered abelian group with finite spines and let  $\{H_i\}_{i \in I}$  be its definable convex subgroups. Then  $G$  is model complete in the language:*

$$\mathcal{L} = \{0, +, -, \leq, (jc_i + H_i)_{j=0,1; i \in I}\},$$

*where  $c_i$  is a representative for the minimal element of the quotient  $G/H_i$  if it is discrete, 0 otherwise.*



# Model completeness with value group with finite spines

## Theorem (D.M.)

*Let  $K$  be an Henselian valued field of mixed characteristic  $(0, p)$ , finite ramification  $e \geq 1$ , and value group  $G$  with finite spines. If the theory of the residue field  $k$  is model complete in the language of rings, then the theory of  $K$  is model complete in the language*

*$\mathcal{L} = \{0, +, \cdot, 1, A_{i,j}\}_{j=0,1;i \in I}$  where  $A_{i,j}$  is a predicate such that*

$$A_{i,j}^K = \{a \in K \mid v(a) = jc_i^G \pmod{H_i}\},$$

*where the  $(H_i)_{i \in I}$  are the definable convex subgroups of  $G$  and  $c_i$  is a representative for the minimal element of the quotient  $G/H_i$  if it is discrete, 0 otherwise.*



# Proof

Assume  $K_1, K_2$   $\aleph_1$ -saturated. Note that:

- $G_1 \preceq G_2$  implies  $G_1 / \langle 1_{G_1} \rangle_{conv} \preceq G_2 / \langle 1_{G_2} \rangle_{conv}$ ;
- AKE in the equicharacteristic 0 case obtained by coarsening holds resplendently considering an expansion of the language of groups;
- the valuation is still  $\exists$  and  $\forall$ -definable by the same formula;
- if  $(H_i)_{i \in I}$  is an enumeration of the definable convex subgroups of  $G$ , it suffices to add predicates  $A_{i,j}$  for  $j = 0, 1$  to  $\mathcal{L}_{ring}$  to have model completeness in a one sorted language.



# Example. Hahn series in one variable

Consider the field of Hahn series  $\mathbb{Q}_p((t^{\mathbb{Z}}))$  and the valuation

$$\text{val} : \mathbb{Q}_p((t^{\mathbb{Z}})) \longrightarrow \mathbb{Z} \times \mathbb{Z}$$

such that

$$O_{\text{val}} = \{x \mid \text{val}(x) \geq 0\} = \{x \mid v_t(x) > 0 \text{ or } v_t(x) = 0 \wedge v_p(\text{ac}_t(x)) \geq 0\}.$$

By the Theorem,  $\text{Th}((\mathbb{Q}_p((t^{\mathbb{Z}})), \text{val}))$  is model complete in the language of rings together with two predicates  $A_0, A_1$  such that

$$\begin{aligned} A_0^{\mathbb{Q}_p((t^{\mathbb{Z}}))} &= \{x \mid \text{val}(x) \in \{0\} + \mathbb{Z}\} \\ &= \left\{x \mid x = \sum_{i \geq 0} a_i t^i, a_0 \neq 0\right\} = \mathbb{Q}_p + (t)^{>0}. \end{aligned}$$

$$\begin{aligned} A_1^{\mathbb{Q}_p((t^{\mathbb{Z}}))} &= \{x \mid \text{val}(x) = (1, 0) \pmod{\{0\} + \mathbb{Z}}\} \\ &= \left\{x \mid x = \sum_{i \geq 1} a_i t^i, a_1 \neq 0\right\} = (t)^{>0}. \end{aligned}$$



# Example. Hahn series with many variables

We can consider the valuation

$$\text{val}_n : \mathbb{Q}_p((t_1^{\mathbb{Z}})) \dots ((t_n^{\mathbb{Z}})) \longrightarrow \bigoplus_{i=1}^{n+1} \mathbb{Z}$$

such that

$$O_{\text{val}_n} = \bigcup_{i=0}^n O^i,$$

where

$$\begin{aligned} O^n &= \{x \mid v_{t_n}(x) > 0\} \\ O^{n-1} &= \{x \mid v_{t_n}(x) = 0 \wedge v_{t_{n-1}}(\text{ac}_{t_n}(x)) > 0\} \\ &\vdots \\ O^0 &= \{x \mid v_{t_n}(x) = 0 \wedge v_{t_{n-1}}(\text{ac}_{t_n}(x)) = 0 \wedge \dots \\ &\quad \wedge v_{t_1}(\text{ac}_{t_2}(\dots(\text{ac}_{t_n}(x)))) = 0 \wedge v_p(\text{ac}_{t_1}(\dots(\text{ac}_{t_n}(x))..)) > 0\} \end{aligned}$$

V:

# Example. Hahn series with many variables

By the Theorem, the theory of the valued field  $\mathcal{K} = (\mathbb{Q}_p((t_1^{\mathbb{Z}})) \dots ((t_n^{\mathbb{Z}})), \text{val}_n)$  is model complete in the language of rings together with predicates  $A_{i,j}$ ,  $i = 1, \dots, n$ ,  $j = 0, 1$ , such that

$$\begin{aligned} A_{i,0}^{\mathcal{K}} &= \{x \in \mathcal{K} \mid \text{val}_n(x) \in H_i\} \\ &= \{x \in \mathcal{K} \mid x \in \mathcal{O}^i \wedge v_{t_i}(\text{ac}_{t_{i+1}}(\dots(\text{ac}_{t_n}(x))\dots)) = 0\}. \end{aligned}$$

$$\begin{aligned} A_{i,1}^{\mathcal{K}} &= \{x \in \mathcal{K} \mid \text{val}_n(x) = c_i^G \pmod{H_i}\} \\ &= \{x \in \mathcal{K} \mid x \in \mathcal{O}^i \wedge v_{t_i}(\text{ac}_{t_{i+1}}(\dots(\text{ac}_{t_n}(x))\dots)) = 1\}. \end{aligned}$$



# Index

## 1 Valued fields

## 2 Transfer principles

- Model completeness for hens vf  $(0,p)$  valued in a Z-group
- Model completeness for hens vf  $(0,p)$  valued in an oag with finite spines
- Model completeness for hens vf  $(0,p)$  valued in an oag with spine of order type  $\omega^*$  and no colors





# The theory of the lexicographic sum of $\mathbb{Z}$

**Example.** Assume that  $(G_\gamma)_{\gamma \in \Gamma}$  is a family of non-trivial archimedean ordered abelian groups, where  $(\Gamma, <)$  is an ordered set. Consider the Hahn product

$$H = \{f \in \prod_{\gamma \in \Gamma} G_\gamma : f \text{ has well ordered support}\}.$$

Then the induced structure on the spine of  $H$  is  $(\Gamma, <, C_\phi)_{\phi \in \mathcal{L}_{oag}}$ , where  $C_\phi$  are unary predicates.



## QE

Let  $\mathcal{L}$  be the language consisting of

- the main sort  $G$  with  $+, -, 0, <, \equiv_m$  ( $m \in \mathbb{N}$ );
- an auxiliary sort  $\Gamma$  with  $<, 0, \infty, s : \Gamma \rightarrow \Gamma$ ;
- $val^n : G \rightarrow \Gamma$  ( $n \in \mathbb{N}, n \neq 1$ ),
- an unary predicate  $=^\bullet k_\bullet$  on  $G$  for each  $k \in \mathbb{Z} \setminus \{0\}$ ,
- an unary predicate  $\equiv_m^\bullet k_\bullet$  on  $G$  for each  $m \geq 2$  and  $k \in \{1, \dots, m-1\}$ .



## QE

## Fact

*Let  $G$  be an oag with spine of order type  $\omega^*$  and no colors. Then the theory of  $G$  has quantifier elimination in  $\mathcal{L}$ , where*

- $\Gamma = \omega^* \cup \{\infty\}$ ,
- $s(n) = n + 1$ ,
- for every  $a \in G$ ,  $\text{val}^n(a) := \text{minsupp}(a \bmod nG)$  if  $a \notin nG$ ,  
 $\text{val}^n(a) := \infty$  otherwise (or equivalently  $\text{val}^n(a)$  is the index  $i$  of the largest convex subgroup  $H_i$  such that  $a \notin H_i + nG$ ),
- for every  $a \in G$ ,  $a \equiv^\bullet k_\bullet$  if  $a + H_i$  is  $k$  times the minimal element of  $G/H_i$  for some  $i \in G$ ,
- for every  $a \in G$ ,  $a \equiv_m^\bullet k_\bullet$  if  $a + H_i$  is congruent modulo  $m$  to  $k$  times the minimal element of  $G/H_i$  for some convex subgroup  $H_i$ .



# Model completeness

## Proposition (D.M.)

*Let  $G$  be an oag with spine of order type  $\omega^*$  and no colors. Then the theory of  $G$  is model complete in the one sorted language  $\mathcal{L}$  consisting of*

- $+, -, 0, <$ ,
- for every  $n, m \in \mathbb{N}$  a relation symbol  $|^{n,m}$ ,
- for every  $n, m \in \mathbb{N}$  a binary predicate  $\bar{s}^{n,m}$ ,
- an unary predicate  $=^{\bullet} 1$ .

where

- $x|^{n,m}y \iff val^n(x) \leq val^m(y)$ ,
- $\bar{s}^{n,m}(x, y) \iff val^m(y) = s(val^n(x))$ ,
- for every  $a \in G$ ,  $a =^{\bullet} 1$  if  $a + H_i$  is the minimal element of  $G/H_i$  for some convex subgroup  $H_i$ .

# Model completeness with value group an oag with spine of order type $\omega^*$ and no colors

## Theorem (D.M.)

Let  $K$  be an Henselian valued field with the same properties and value group an oag  $G$  with spine of order type  $\omega^*$  and no colors. If  $\text{Th}(k)$  is m.c. in  $\mathcal{L}_{\text{rings}}$ , then  $\text{Th}(K)$  is m.c. in  $\mathcal{L}_{\text{rings}}$  together with

- for every  $n, m \in \mathbb{N}$  a relation symbol  $\|^{n,m}$ ,
- for every  $n, m \in \mathbb{N}$  a binary predicate  $\$^{n,m}$ ,
- an unary predicate  $A$ ,

where

- for every  $x, y \in K$ ,  $x\|^{n,m}y \iff \text{val}^n(v(x)) \leq \text{val}^m(v(y))$ ,
- for every  $x, y \in K$ ,  $\$(x, y) \iff \text{val}^m(v(y)) = s(\text{val}^n(v(x)))$ ,
- $A^K = \{x \in K \mid v(x) = \bullet 1_\bullet\}$ .

# Example infinite many variables

This language gives model completeness for the following valued field. Consider the field of Hahn series over  $\mathbb{Q}_p$  in infinitely many indeterminates

$$\mathcal{K} = \bigcup_{n \in \mathbb{N}} \mathbb{Q}_p((t_1^{\mathbb{Z}})) \dots ((t_n^{\mathbb{Z}})).$$

We can define, from the valuations  $val_n$  over  $\mathbb{Q}_p((t_1^{\mathbb{Z}})) \dots ((t_n^{\mathbb{Z}}))$ , a valuation  $val_\infty$  over  $\mathcal{K}$  with values in  $\bigoplus_{i < \omega^*} \mathbb{Z}$ , such that

$$O_{val_\infty} = \bigcup_{n \in \mathbb{N}} O_{val_n}.$$

